





Diana Stralberg, Mark Herzog, Nils Warnock, Nadav Nur


San Francisco Bay Estuary

- Designated site of hemispheric importance for shorebirds
- Important waterfowl wintering area
- Largest area of tidal marsh habitat on west coast for breeding songbirds and rails

 South SF Bay waterbird use particularly high




Historic Habitat Change: Diking, Filling, Dredging




Current Habitat Change: Salt Pond Restoration

- 5,471 ha of salt ponds purchased by state and federal agencies for restoration to tidal and managed marsh
- California
 Coastal
 Conservancy
 overseeing
 restoration






Range of Possible Habitats



Model Objectives

- Identify bird-related costs & benefits of converting salt ponds to tidal marsh
- Develop quantitative ability to help guide:
 - Design of tidal marsh restoration projects
 - Management of remaining salt ponds
 - Optimal configuration of tidal marshes and salt ponds


Study Areas



Survey Methods



Site Characteristics



Salt Ponds


- Salinity (ppt)
- Pond size



Ponded area

Landscape Characteristics

- Distance to open bay, development
- Surrounding landscape composition



Tidal Marsh Model Parameters*

	Songbirds and Rails	Small Shorebirds	Large Shorebirds	Dabbling Ducks	Diving Ducks
Site Characteristics					
Large Channel (> 8 m) Area			+	+	+
Ponded Area	-	+	+	+	
Landscape Characteristics					
Salt Ponds within 1 km	-				
Mudflat within 1 km		+		+	
Marsh within 1 km	+				
Open Bay within 1 km					+


^{*} Signs of coefficients from multiple regression models (bird variables log-transformed)

Site-level Restoration Scenarios*

1 - Salt ponds retained at current salinities (no change)

All salt ponds restored to tidal marsh:

2: minimum large channels and ponded areas



3: mean large channels and ponded areas




4: maximum large channels and ponded areas




^{*} Landscape conditions incorporated into models.

Comparing Restoration Scenarios




Western Sandpiper

Scenario Comparisons, Small Shorebirds





Scenario Comparisons, Diving Ducks



Conclusions from Phase I

- Loss of salt ponds may cause substantial reduction in waterbird numbers, especially small shorebirds and diving ducks.
- Potential to reduce and/or avoid waterbird losses through design and management of individual restoration sites.
- Critical for waterbirds to retain some salt ponds in a habitat mosaic (more important than tidal marsh design and management).

Recommendations from Phase I

- A few well-managed salt ponds can go a long way for waterbirds.
- Trade-offs are inevitable try to optimize, but can't treat all species equally.
- A diverse wetland mosaic (salt ponds, tidal marsh, mudflat) will provide the best chance for all birds.
- Restoration should proceed with caution; monitor results along the way.

Acknowledgements

Funders:

California Coastal Conservancy
Gabilan Foundation
Mary A. Crocker Trust
Rintels Charitable Trust
Tides Foundation

Partners:

San Francisco Bay Bird Observatory
U.S. Geological Survey
U.S. Fish & Wildlife Service
(Don Edwards San Francisco Bay NWR)



Future Monitoring and Research Needs

- Long-term monitoring program
 - document population trends and habitat requirements
 - inform adaptive management
- Food resources
 - Salton Sea (fish, invertebrates)
 - created habitats (invertebrates)
- Disease and contaminants
- Important habitat features (e.g., nesting sites)
- Agricultural habitats